

TMUP-HEL-9406

May, 1994

Solitons in Nonlinear Schrödinger Model and the Collective Ground State of One-Dimensional Delta-Function Gas

Hideaki Hiro-Oka^{*†} and Hisakazu Minakata[‡]

*Department of Physics
Tokyo Metropolitan University
1-1 Minami-Osawa Hachioji, Tokyo 192-03, Japan*

Abstract

We examine one-dimensional Bose gas interacting with delta-function potential using the large- N collective field theory. We show that in the case of attractive potential the uniform ground state is unstable to small perturbations and the instability is cured by formation of a collective ground state, “bright soliton” configuration in corresponding nonlinear Schrödinger field theory.

^{*}Research Fellow of Japan Society for the Promotion of Science

[†]e-mail: hiro-oka@phys.metro-u.ac.jp

[‡]e-mail: minakata@phys.metro-u.ac.jp

The large- N collective field theory formulated by Jevicki and Sakita [1][2] has been known as a powerful nonperturbative tool in dealing with many-body collective phenomena. It is remarkable that its applicability extends over wide range of problems such as, plasma oscillations [1][2], matrix models [1]-[3], the quantum Hall effect [4], and anyons [5].

In this paper we demonstrate that the large- N collective field theory is a really powerful machinery in uncovering collective modes at high densities. We shall do this in a particular model of bosons in 1 dimension interacting via a delta-function potential. Despite that our discussion will be restricted to this particular model we believe that the power of the formalism in hunting collective excitations extends over a wide class of models.

One-dimensional Bose gas with delta-function potential have been known to be soluble [6] since long time ago. Nonetheless, it provides a nontrivial testing ground in the context of the collective field formalism and we will learn some meaningful lessons. First of all we will verify that the formalism reproduces the results obtained by Lieb and Liniger [6] in high-density limit. We observe, not surprisingly, that the ground state energy in the case of repulsive potential is of the order of $\hbar^0$ and is in agreement with that of ref.[6]. We also obtain a classical sound velocity consistent with the classical ground state energy. More importantly, we clarify how the instability of the uniform ground state in the case of attractive potential can be stabilized by forming a collective ground state, a “bright soliton” configuration of the corresponding nonlinear Schrödinger field theory. We also give some remarks on the role of “dark soliton” configurations in the case of repulsive potential.

The system of N particles interacting with a delta-function potential is described by the Hamiltonian

$$H = -\frac{\hbar^2}{2m} \sum_{a=1}^N \frac{\partial^2}{\partial x_a^2} + 2c \sum_{a>b} \delta(x_a - x_b). \quad (1)$$

Our notation is the same as that of Lieb and Liniger [6] except that they take $m = 1/2$. Also we shall explicitly denote $\hbar$ throughout this paper for convenience of our discussions. The canonical approach to the problem is, of course, to solve the eigenvalue problem with the Hamiltonian (1) using a symmetric (Bose) and an antisymmetric (Fermi) wave functions. This is beautifully done by Lieb and Liniger in the former case by using the Bethe ansatz technique.

Instead, we shall develop a collective field theory treatment of this system in this paper. Following the well-established way [2] we rewrite the Hamiltonian (1) in terms of the density variables

$$\phi(x) = \frac{1}{N} \sum_{a=1}^N \delta(x - x_a) \quad (2)$$

and its canonical conjugate $\pi(x)$. Here the density variable is normalized such that

$$\int dx \phi(x) = 1. \quad (3)$$

The comments that follow are well-known but are in order for readers who are unfamiliar with the large- N collective field theory: One should notice that the Hamiltonian we shall work with is not the one which we will obtain by just changing the variables from x_a to $\phi(x)$. Let us denote the Jacobian as J of the transformation of x_a to $\phi(x)$ in (2), or more precisely speaking, to its Fourier transformation

$$\begin{aligned} \phi_k &= \int dx e^{-ikx} \phi(x) \\ &= \frac{1}{N} \sum_{a=1}^N e^{-ikx_a}. \end{aligned} \quad (4)$$

Then the functional space we are to work with is that of Schrödinger's wave function $J^{1/2}\Psi$ not Ψ , where Ψ denotes the Schrödinger's wave function of the original Hamiltonian (1). Therefore, the Hamiltonian for our system written in terms of $\phi(x)$ is $J^{1/2}H(\phi, \pi)J^{-1/2}$, and this is the Hamiltonian which possesses hermiticity in the functional space of $J^{1/2}\Psi$. We shall denote this new Hamiltonian as H_{eff} .

We obtain H_{eff} in the following form:

$$\begin{aligned} H_{eff} &= \frac{\hbar^2}{2mN} \int dx (\partial\pi) \phi (\partial\pi) + \frac{\hbar^2 N}{8m} \int dx \frac{(\partial\phi)^2}{\phi} \\ &\quad + N^2 c \int dx \phi^2 - \lambda N \left(\int dx \phi - 1 \right), \end{aligned} \quad (5)$$

where ∂ is an abbreviation of $\frac{\partial}{\partial x}$. This is the Hamiltonian which we analyze using the large- N technology. Note that we only have a nontrivial interacting

theory under the large- N limit in which cN is finite and $N \rightarrow \infty$. In this case all the terms in (5) are of order N assuming that π is of order N .

Let us discuss the ground state and the low-lying excited states of the system (5). Since the Hamiltonian is of order N we may allow to use saddle-point approximation, e.g., in the path integral formalism. As a result we have the classical Hamilton's equations. For time-independent ground-state configuration they read,

$$\partial(\phi\partial\pi) = 0, \quad (6)$$

$$\frac{\hbar^2}{2m}(\partial\pi)^2 - \frac{\hbar^2}{4m}\frac{\partial^2\phi}{\phi} + \frac{\hbar^2}{8m}\left(\frac{\partial\phi}{\phi}\right)^2 + 2Nc\phi - \lambda = 0. \quad (7)$$

We require that the ground state is translationally invariant. Then the solution is, rather trivially,

$$\phi_0(x) = \frac{1}{L}, \quad \pi_0(x) = 0, \quad (8)$$

where we have put the system into the box of length L . By virtue of (7) we have the relation between the chemical potential λ and the density of particles $\rho = N/L$ as $\lambda = 2c\rho$. Using (5) and (8) we obtain the ground state energy of the system as

$$\begin{aligned} E_0 &= N^2 c \cdot \frac{1}{L}, \\ &= Nc\rho, \end{aligned} \quad (9)$$

in agreement with the result of Lieb-Liniger [6] at high-density ($\rho \rightarrow \infty$) limit in the repulsive potential $c > 0$. Thus our machinery works in the large- N limit. We note that the ground state energy (9) is a classical one, namely, it is free from $\hbar$. This feature is shared by the exact result obtained in ref.[6] as one can verify by simple dimensional analysis. While somewhat curious at first site it is an example of the well-known fact that the system becomes classical in high density limit.

So far our discussion is valid not only for repulsive ($c > 0$) potential but also for attractive ($c < 0$) potential. Apparently the ground state (8) exists also for the latter case in contradiction to the claim given in ref.[6].

To resolve the problem and to verify the validity of the large- N collective field theory we examine low-lying excitations as small fluctuations around the

ground state (8). We employ plane-wave basis to expand the fluctuations as

$$\begin{aligned}\phi(x) &= \frac{1}{L} \left(1 + \sum_k \sqrt{\frac{k^2}{N}} e^{ikx} Q_k \right), \\ \pi(x) &= \frac{1}{\hbar} \sum_k \sqrt{\frac{N}{k^2}} e^{-ikx} P_k.\end{aligned}\tag{10}$$

Inserting (10) into the Hamiltonian (5) and picking up the order N^0 term [order $N^{1/2}$ term cancels because of equations of motion (6) and (7)] we obtain the Hamiltonian of harmonic oscillator form

$$H_{N^0} = \sum_k \left(\frac{1}{2m} |P_k|^2 + \frac{m}{2} \omega^2(k) |Q_k|^2 \right),\tag{11}$$

where

$$\hbar^2 \omega^2(k) = \left(\frac{\hbar^2}{2m} k^2 \right)^2 + \frac{2c\rho}{m} \hbar^2 k^2.\tag{12}$$

In the case of repulsive potential each term of (12) allows a simple physical interpretation: the first term is an obvious kinetic energy term, and the second one is the energy of sound wave with the velocity

$$v_s = \sqrt{\frac{2c\rho}{m}}.\tag{13}$$

By a well-known macroscopic argument the sound velocity is given by

$$v_s = \sqrt{-\frac{L}{m\rho} \frac{\partial P}{\partial L}}\tag{14}$$

where P is the pressure,

$$P = -\frac{\partial E_0}{\partial L}.\tag{15}$$

It is amusing to observe that the use of (14), (15), and (9) reproduces the expression (13) for the sound velocity. This is, of course, nothing but repeating the argument in ref.[6] under the high-density approximation. We just want to point out that the sound velocity (13) has a classical value because the ground state energy (9) is classical. Thus we have seen that the large- N collective field theory works for Bose gas with repulsive delta-function potential in the high-density limit.

In the case of attractive potential, $c < 0$, the interpretation of (12) is quite different. At sufficiently small k the frequency ω becomes imaginary signaling the instability of the ground state (8). In this way we have reached a conclusion that there is no stable *uniform* ground state for attractive potential, in agreement with ref.[6].

Now we turn to the discussion of collective ground state. Namely, we show that there exist solutions in (6) and (7) which can be interpreted as collective excitation of Bose particles at high-densities.

To this goal we introduce a Schrödinger variable

$$\psi(x) = \sqrt{\phi(x)} e^{-i\pi(x)}. \quad (16)$$

Notice that there is no direct connection between this ψ and the many-body Schrödinger wave function of the original problem. Then it is easy to show that the equations (6) and (7) can be rewritten as

$$-\frac{\hbar^2}{2m} \frac{d^2}{dx^2} \psi(x) + 2Nc|\psi|^2 \psi(x) = \lambda \psi(x) \quad (17)$$

This is nothing but the classical nonlinear Schrödinger equation.

It is well known that the equation (17) admits an infinite number of soliton solutions [7]. Solitons in theories with attractive and repulsive interactions are called “bright” and “dark” solitons, respectively. These names originate in its application to the problem of nonlinear optical pulses in dispersive dielectric fibers [8].

As discussed before there exists no stable uniform ground state for $c < 0$ case. However, we see that the collective ground state can exist, which corresponds to the bright soliton. In the case of repulsive interaction we can construct the dark soliton solution besides the uniform ground state.

For the purpose of illuminating the general picture we work with a simple ansatz of the bright soliton solution of the form

$$\psi(x) \propto \text{sech } ax. \quad (18)$$

In our language we assume

$$\phi = A \text{sech}^2 ax, \quad \pi = B, \quad (19)$$

from the relation (16). The unknown constants a , A , and B are determined by the requirement that (19) are solutions to (6) and (7). Substituting (19) into (6) and (7) we obtain

$$\left(\frac{\hbar^2 a^2}{m} + 2NcA\right) \operatorname{sech}^2 ax - \left(\lambda + \frac{\hbar^2 a^2}{2m}\right) = 0. \quad (20)$$

This relation lead to

$$A = -\frac{\hbar^2 a^2}{2Ncm}, \quad \lambda = -\frac{\hbar^2 a^2}{2m}, \quad (21)$$

apart from a trivial case $a = \lambda = 0$. By definition ϕ and A must be positive, so we have two cases: i) $c < 0$ and $a \in \mathbb{R}$, ii) $c > 0$ and $a = i\alpha$, $\alpha \in \mathbb{R}$. As can be seen the case ii) yields the unphysical solution with singularities. Thus we have

$$\phi = \frac{a^2}{2N\tilde{c}m} \operatorname{sech}^2 ax, \quad (22)$$

where $\tilde{c} = -c > 0$. On the other hand π is an arbitrary constant. Due to the normalization condition for ϕ , a is determined as

$$a = \frac{N\tilde{c}m}{\hbar^2}. \quad (23)$$

This is the collective ground state. We obtain the ground state energy as

$$E_0 = -\frac{m}{6\hbar^2} N(Nc)^2, \quad (24)$$

by inserting (22) to the effective Hamiltonian. Note that this energy is in agreement with that of a static bright soliton solution of the classical non-linear Schrödinger problem.

Next we consider the dark soliton solution of the form

$$\phi = A \tanh^2 ax, \quad \pi = B, \quad (25)$$

in the similar way. Note that this is a special case of generic dark soliton solutions. Substituting (25) into (6) and (7) we obtain

$$\left(2NcA - \frac{\hbar^2 a^2}{m}\right) \tanh^2 ax + \left(\frac{\hbar^2 a^2}{m} - \lambda\right) = 0. \quad (26)$$

By the same discussion we have

$$A = \frac{\hbar^2 a^2}{2Ncm}, \quad \lambda = \frac{\hbar^2 a^2}{m}. \quad (27)$$

The constant a is given by the normalization of ϕ ,

$$|a| = \frac{1}{L} \left(1 + \sqrt{1 + \frac{2Ncm}{\hbar^2} L} \right). \quad (28)$$

The ground state energy is given as

$$E_0 = N\rho c + \frac{4}{3} \frac{\hbar\rho}{m} \sqrt{2m\rho c} \quad (29)$$

apart from the correction of the order of e^{-L} . The first term of (29) is nothing but the energy of the uniform ground state while the second term is the quantum correction of order $\hbar$.

To summarize we make a few remarks:

i) We have clarified, within the framework of large- N collective field theory, how the instability of the uniform ground state of the attractive delta-function gas can be cured by the formation of the collective ground state. The feature that the ground state itself is a non-uniform collective state seems to be very new to us.

ii) In the case of repulsive potential the ground state is uniform and is stable against small perturbations. We have reproduced Lieb-Liniger's result on the energy of the fluctuation mode in high-density limit. We also have dark soliton excitation on this ground state.

iii) The quantum meaning of the bright soliton solution of the classical nonlinear Schrödinger field theory has been thoroughly discussed by Wadati and coworkers [9]. The large- N collective field formalism can be viewed as a way of constructing quantum field theory starting from N -body Schrödinger problem. This is achieved by taking the thermodynamic limit, $N \rightarrow \infty$ keeping N/L fixed. Whereas we have discussed the high-density limit, $N \rightarrow \infty$ keeping L large but fixed in this paper and the clear relationship with classical soliton solutions has emerged. Therefore, the large- N collective theory provides an another way of understanding quantum correspondence with the classical object.

Works of H. M. and H. H. are partially supported by Grant-in-Aid for Scientific Research of the Ministry of Education, Science and Culture #05640355 and #H4-2028, respectively.

References

- [1] A. Jevicki and B. Sakita, *Nucl. Phys.* **B165** (1980) 511, **B185** (1981) 89.
- [2] B. Sakita, Quantum Theory of Many-Variable Systems and Fields (World Scientific, Singapore, 1985).
- [3] I. Andrić, A. Jevicki, and H. Levine, *Nucl Phys.* **B215** (1983) 307.
- [4] B. Sakita, D.-N. Sheng, and Z.-B. Su, *Phys. Rev.* **B44** (1991) 11510; B. Sakita and Z.-B. Su, *Rochester 40 Years* (1990) 179.
- [5] H. Hiro-Oka and H. Minakata, *Mod. Phys. Lett.* **A8** (1993) 3547.
- [6] E. H. Lieb and W. Liniger, *Phys. Rev.* **130** (1963) 1605.
- [7] V. E. Zakharov and A. B. Shabat, *Sov. Phys. JETP* **34** (1972) 62; *Sov. Phys. JETP* **37** (1973) 823.
- [8] A. Hasegawa and F. Tappert, *Appl. Phys. Lett.* **23** (1973) 142; *Appl. Phys. Lett.* **23** (1973) 171.
- [9] M. Wadachi and M.-A. Sakagami, *J. Phys. Soc. Jpn* **53** (1984) 1933; M. Wadati, A. Kuniba, and T. Konishi, *J. Phys. Soc. Jpn* **54** (1985) 1710.